

5. Phillips EJ, Keystone JS, Kain KC. Failure of combined chloroquine and high-dose primaquine therapy for *Plasmodium vivax* malaria acquired in Guyana, South America. *Clin Infect Dis*. 1996;23:1171–3.
6. Ruebush TK II, Zegarra J, Cairo EM, Andersen M, Green DR., Pillai W, et al. Chloroquine-resistant *Plasmodium vivax* malaria in Peru. *Am J Trop Med Hyg*. 2003;69:548–52.
7. Alecrim MGC. Estudo clínico, resistência e polimorfismo parasitário na malária pelo *Plasmodium vivax*, Manaus – AM. Brasília: UNB, 2000. PhD Thesis, Faculdade de Medicina, Núcleo de Medicina Tropical, Universidade de Brasília, 2000.
8. Baird JK, Leksana B, Masbar S, Fryauff DJ, Sutanihardja MA, Suradi FS, et al. Diagnosis of resistance to chloroquine by *Plasmodium vivax*: timing of recurrence and whole blood levels. *Am J Trop Med Hyg*. 1997;56:621–6.
9. Pan-American Health Organization. Generic protocol for antimalarial drug-efficacy studies in the Américas. Efficacy of chloroquine for the treatment of *Plasmodium vivax* malaria. 2004. [cited 2007 May 15]. Available from <http://www.paho.org/english/AD/DPC/CD/mal-antimalarials.htm>
10. Yonemitsu K, Koreeda A, Kibayashi K, Ng'walali P, Mbonde M, Kitinya J, et al. HPLC analysis of anti-malaria agent, chloroquine in blood and tissue from forensic autopsy cases in Tanzania. *Leg Med (Tokyo)*. 2005;7:113–6.

Address for correspondence: Franklin Simoes de Santana Filho, The Foundation for Tropical Medicine of Amazonas, Av Pedro Teixeira 25 Planalto, Manaus, Amazonas 69040000, Brazil; email: fsimoes@fmt.am.gov.br

Avian Influenza Risk Perceptions, Laos

To the Editor: After the 2004 outbreak of highly pathogenic avian influenza (HPAI) in poultry in Lao People's Democratic Republic (PDR), the Ministry of Health implemented extensive virologic surveillance (1,2). Surveillance began in July 2005, and

by early 2006, only sporadic cases were found. In July 2006, an outbreak of HPAI was confirmed on 2 chicken farms in Vientiane, the capital city of Lao PDR (1,3). Most of Laos' ≈20 million chickens are kept on family-owned backyard farms; 3.2 million are on commercial farms (4). This production meets 80% of Lao poultry (chicken, duck, goose, quail) needs; imports from neighboring countries, either through legal trade or cross-border smuggling, account for the rest (3). Common poultry diseases occur frequently during the cold season, and lack of reporting of poultry deaths is of concern (4).

Until February 2007, no human cases of influenza A (H5N1) had been reported in Lao PDR. To learn more about Laotians' knowledge of HPAI and perceptions of their risk, we conducted a cross-sectional survey.

In March–April 2006, participants in 3 settings (Vientiane, urban; Oudomxay, semiurban; Attapeu Province and Hinheub District, both rural) were interviewed in the Lao language by means of a standardized 33-question survey. We recorded information about behavior, poultry handling and keeping practices, and poultry deaths. We used multivariate analysis (Stata, version 8; Stata Corporation, College Station, TX, USA) to analyze the factors associated with behavior changes.

Using a random sampling list of visitors and vendors, we interviewed 461 respondents in 4 Vientiane city markets (Vientiane has 114,793 households and 3,700 registered poultry farms) (5). Semiurban respondents were recruited in Oudomxay (40,987 households, 715 poultry farms), an active trading zone near the Chinese border. Rural respondents were recruited from Hinheup District and in Attapeu (19,050 households, 360 poultry farms), near the Vietnam border. Twenty villages were randomly selected, and 10 participants per village were randomly selected for interview. Approval for the investigation was

obtained from the health and market authorities. Oral consent for interview was obtained from participants.

A total of 842 participants were interviewed (Table). Differences in occupation and literacy were associated with different study areas. Differences in participant sex and age were also noted because, in the rural areas, interviews took place in the home. A total of 583 (69.3%) participants were female: 302 (65.5%), 139 (68.2%), and 150 (79.3%), in urban, semiurban, and rural areas, respectively; $p = 0.002$, 95% confidence interval 66–72. Mean ages for participants in these areas were 41 (range 40–43), 34 (range 32–36), and 38 (range 37–41) years, respectively; $p < 0.001$. Animal breeding was conducted by 50% of families. Daily close exposure to poultry was common (39.6%). Few families owned a henhouse, and no special handling of poultry was reported. Rates of poultry vaccination against common poultry diseases were higher in urban and semiurban areas; veterinary surveillance was low (10.2%).

Overall, 96.9% of respondents had already heard of HPAI, mainly through television. Urban residents ranked it as the most well-known poultry disease, but rural residents ranked it fifth. Less than half of the respondents had some knowledge of the disease signs and symptoms for humans and poultry; 28.4% could describe 1 symptom. Half of the respondents believed that they were not at risk for human avian influenza or that their poultry were not at risk for it. Respondents in urban and semiurban areas knew more about avian influenza than those in rural areas.

During the cold season, poultry deaths were higher in the north (colder) and south than in Vientiane. The poultry mortality rate during the cold season was similar to that of Cambodia (6). Behavior regarding poultry deaths differed between areas. Despite a high rate of poultry deaths, none of the interviewees had notified authorities. Since hearing about HPAI, 67.1%

Table. Avian influenza knowledge, risk perception, and poultry-keeping behavior, Lao People's Democratic Republic*

Characteristic	Urban, n (%)	Semiurban, n (%)	Rural, n (%)	Total, n (%)	p value	95% CI
Total persons interviewed	461	192	189	842		
Illiterate	175 (37.9)	60 (31.2)	181 (95.7)	416 (49.4)	<0.001	47.1–54
Occupation						
Housewife	126 (27.3)	24 (12.5)	94 (50)	244 (28.9)	<0.001	32–25.9
Farmer	25 (5.4)	36 (18.75)	75 (40)	136 (16.1)	<0.001	13.7–18.6
Government worker	103 (22.4)	22 (11.5)	3 (1.5)	128 (15.2)	<0.001	12.8–17.6
None	2 (4.3)	0	24 (12.6)	36 (4.2)	<0.001	2.9–5.6
Keep poultry	185 (40.2)	97 (50.5)	159 (84.3)	441 (59.4)	<0.001	19 (17–20)
≥1 poultry death, past 2 mo†	58 (31.3)	84 (86.5)	95 (59.7)	239 (54.1)	<0.000	49.5–58.8
Any poultry deaths, past 2 y	95 (51.3)	62 (63.9)	141 (88.6)	298 (65.5)	<0.001	63.2–71.9
Response to dead poultry (n = 399)‡						
Bury dead chickens	105 (56.7)	87 (89.6)	118 (74.2)	310 (70.2)	<0.001	66–74.6
Throw out dead chickens	50 (27.0)	5 (5.1)	9 (5.6)	64 (14.5)	<0.001	11.2–17.8
Eat dead chickens	1 (0.5)	2 (2.0)	7 (4.4)	10 (2.2)	0.06	0.9–3.7
Treat other chickens	0	0	5 (2.6)	5 (0.5)	<0.001	0.07–1.1
Apply lime to backyard	0	8 (1.7)	1 (0.5)	9 (1.0)	<0.001	0.03–1.7
Sell dead chickens	0	1 (1.0)	0	1 (0.1)	0.1	0.00–0.3
Report dead chickens	0	0	0	0	NA	NA
Poultry location						
Henhouse	39 (21.0)	4 (4.4)	7 (4.4)	50 (11.3)	<0.001	8.4–14.3
Inside house	8 (4.3)	1 (1.03)	2 (12.6)	11 (2.4)	0.003	1–3.9
Near house (<5 m)	78 (42.2)	59 (61)	28 (17.7)	165 (37.4)	<0.001	32.9–41.9
Far from house (>5 m)	58 (31.3)	30 (31)	114 (71.7)	202 (45.8)	<0.001	41.2–50.5
Regular poultry vaccination	81 (43.7)	54 (55.6)	19 (11.9)	154 (34.2)	<0.001	30.5–39.4
Information source						
Never heard	8 (1.7)	11 (5.1)	7 (3.7)	26/837 (3.1)	0.02	1.9–4.3
Heard from television	388 (86.4)	158 (87.8)	178 (97.8)	724 (89.2)	<0.001	(86.4–90.8)
Heard from radio	19 (4.2)	12 (6.6)	4 (2.2)	35 (4.3)	0.1	(3.02–5.9)
Read in paper	6 (1.3)	1 (0.5)	0	7 (0.8)	0.003	(0.34–1.8)
Perceive risk for avian influenza						
In Laos	369 (81.6)	110 (60.7)	8 (4.3)	487 (59.6)	<0.001	56.3–63
At home	293 (64.8)	72 (40.0)	5 (2.6)	370 (45.7)	<0.001	41.9–48.8
Unable to describe human disease	116 (25.6)	116 (63.7)	182 (97.5)	414 (50.7)	<0.001	47.3–54.2
Able to describe as lethal for poultry	306 (67.5)	90 (49.7)	2 (1.0)	398 (48.7)	<0.0001	45.3–52.2
Behavior change‡	416 (91.8)	125 (69.0)	7 (3.8)	548 (67.1)	<0.0001	63.9–70.4
Stopped eating chicken	328 (72.4)	120 (66.2)	0	448 (54.9)	<0.000	51.5–58.3
Avoided contact	348 (76.8)	60 (33.1)	3 (1.6)	411 (50.3)	<0.000	46.9–53.8
Stopped keeping poultry	335 (73.9)	13 (7.1)	1 (0.5)	349 (42.7)	<0.000	39.4–46.2
Wore mask	338 (74.6)	10 (5.5)	1 (0.5)	349 (42.7)	<0.000	39.4–46.2
Washed hands after contact	100 (22.0)	3 (1.6)	1 (0.5)	104 (12.7)	<0.002	10.5–15
Ate well-cooked chicken	155 (34.2)	3 (1.6)	1 (0.5)	159 (19.4)	<0.000	16.8–22.2

*CI, confidence interval; NA, not applicable.

†Mean nos. of poultry deaths were 15 (range 10–19), 27 (range 22–32), and 15 (range 13–18) for urban, semiurban, and rural areas, respectively. Total mean = 19.3; p<0.0001; 95% CI, 17.0–18.4.

‡95% CIs were 89–94, 62–76, and 1–7 for urban, semiurban, and rural areas, respectively.

respondents, mainly in Vientiane, claimed that they had changed behavior regarding poultry. Multivariate analysis showed the following factors to be associated with behavior change: level of education ($p = 0.002$), urban living ($p < 0.001$), knowledge of avian influenza risk ($p < 0.001$) and disease ($p < 0.001$), owning poultry ($p < 0.001$), and being a government worker ($p < 0.001$).

This study had limitations but provides new insights on Laotians' knowledge and poultry practices with regard to HPAI. Despite a high level of awareness, populations underestimated the risk, particularly those in rural areas. Most respondents were unaware of appropriate poultry-handling measures to reduce risk (6). The claimed changes were higher (more frequent and more substantial) in ur-

ban (91.8%) than in rural sites (3.8%, $p < 0.001$), higher than changes made by their counterparts in Thailand (7), and confirmed by reports after the 2004 outbreaks (8,9). These differences between urban and rural areas might be explained not only by participant characteristics but also by a lower extent of the awareness campaign in rural areas.

Failure to report poultry deaths should be addressed and has several possible explanations. Farmers are accustomed to common yearly poultry deaths, which are not reported. In the absence of an official compensation statement, farmers may fear income loss from massive poultry culling.

Our results emphasize the need for more accurate information about transmission risks, notification requirements, safer behavior and practices, and compensation for losses. Focus also needs to be placed on building capacity in the veterinary system (10). These issues should be integrated in the Laos National Avian Influenza Control and Pandemic Preparedness Plan (2006–2010).

Acknowledgments

We thank the seventh-class students of the Institut Francophone pour la Médecine Tropicale, Vientiane; Lao PDR; and Lao Health Authorities. We thank Evelyne Franon, Vicky Houssiere, Ricarda Monty, and Louise Pelletier for advice and documentation and Philippe Brosman, Cindy Chu, and Ralph Osterwoltdt for revising the final draft.

This study was part of a Master in Tropical Medicine training program of the Institut Francophone pour la Médecine Tropicale, Vientiane, Lao PDR, granted by Agence Universitaire de la Francophonie.

**Hubert Barennes,*
Bertrand Martinez-Aussel,*
Phengta Vongphrachanh,*†
and Michel Strobel***

*Institut Francophone pour la Médecine Tropicale, Vientiane, Lao PDR; and †Ministry of Health, Vientiane, Lao PDR

References

1. Boltz DA, Douangngneun B, Sinthasack S, Phommachanh P, Rolston S, Chen H, et al. H5N1 influenza viruses in Lao People's Democratic Republic. *Emerg Infect Dis.* 2006;10:1593–4.
2. Witt CJ, Malone JL. A veterinarian's experience of the spring 2004 avian influenza outbreak in Laos. *Lancet Infect Dis.* 2005;5:143–5.
3. World Health Organization. Avian influenza update no. 56: 15 August 2006. [cited 2007 Apr 18]. Available from <http://www.wpro.who.int/NR/rdonlyres/13C8C409-2438-4426-A5E3-7339564D1731/0/AIWeekly56WPRO.pdf>
4. Food and Agriculture Organization. Epidemiology of H5N1 influenza in Asia and implications for regional control. [cited 2007 Apr 18]. Available from <http://www.fao.org/ag/againfo/subjects/documents/ai/HPAI-Masseyreport.pdf>
5. National Statistical Centre. Lao statistical yearbook 2003. Vientiane (Lao PDR): Committee for Planning and Cooperation; 2004.
6. Vong S, Coghlan B, Mardy S, Holl D, Seng H, Ly S, et al. Low frequency of poultry-to-human H5N1 virus transmission, southern Cambodia, 2005. *Emerg Infect Dis.* 2006;10:1542–8.
7. Takeuchi MT. Avian influenza risk communication, Thailand. *Emerg Infect Dis.* 2006;12:1172–3.
8. Food and Agriculture Organization. Live-stock report 2006. Rome, 2006. [cited 2007 Apr 18]. Available from <http://www.fao.org/docrep/009/a0255e/a0255e00.htm>
9. US Department of Agriculture. Laos poultry and products: avian influenza 2005. Report no. LA5001. [cited 2007 Apr 18]. Available from <http://www.fas.usda.gov/gainfiles/200503/146119131.doc>
10. World Organization for Animal Health. Laos 2002, veterinarians and technical personnel. [cited 2007 Apr 18]. Available from http://www.oie.int/hs2/gi_veto_pays.asp?c_pays=106&annee=2002

Address for correspondence: Hubert Barennes, Institut Francophone pour la Médecine Tropicale, BP 9516 Vientiane, Lao PDR; email: hubert.barennes@auf.org

Letters

Letters commenting on recent articles as well as letters reporting cases, outbreaks, or original research are welcome. Letters commenting on articles should contain no more than 300 words and 5 references; they are more likely to be published if submitted within 4 weeks of the original article's publication. Letters reporting cases, outbreaks, or original research should contain no more than 800 words and 10 references. They may have one Figure or Table and should not be divided into sections. All letters should contain material not previously published and include a word count.

Norovirus GII.4 Strains and Outbreaks, Australia

To the Editor: Viral gastroenteritis affects millions of people of all ages worldwide, and some seasonality has been observed in outbreak occurrences (1–3). During early 2006 in New South Wales (NSW), a marked increase in outbreaks of gastroenteritis occurred (Figure): 155 outbreaks were reported during the first 5 months compared with 88 outbreaks during 2005. During the first 5 months of 2006, the Enteric Pathogens Laboratory–South Eastern Area Laboratory Services (EPL-SEALS) recorded an increase in norovirus in stool samples, detected by using an enzyme immunoassay (IDEIA Norovirus, DakoCytomation, Cambridgeshire, UK). From January through May 2006, the proportion of samples positive for norovirus increased successively: 0/47 (0%), 1/73 (1.4%), 5/169 (3.0%), 8/106 (7.5%), and 93/413 (22.5%). This trend followed the increasing reports of outbreaks made to the NSW Department of Health (Figure). In May, the rate of norovirus detection (22.5%) was significantly greater than that of any other pathogen (Fisher exact test, $p < 0.0001$), including intestinal parasites, foodborne bacterial pathogens (*Salmonella*, *Shigella*, and *Campylobacter*), and enteric viruses (rotavirus, adenovirus, and astrovirus).

In April 2006, the NSW Department of Health Public Health Real-time Emergency Department Surveillance System (PHREDSS) detected a significant increase in visits for gastroenteritis. This system records cases in real time for each visit to an emergency department from patient demographic information and syndromes diagnosed according to the International Classification of Diseases, version 9, Clinical Modification (ICD-9-CM) (4). Information collected came

Copyright of Emerging Infectious Diseases is the property of Centers for Disease Control & Prevention (CDC) and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.